

Zeroes of holomorphic functions with almost-periodic modulus

Favorov S.Ju.

Abstract

We give necessary and sufficient conditions for a divisor in a tube domain to be the divisor of a holomorphic function with almost-periodic modulus.

AMS classification: 42A75 (32A60, 32A18)

Keywords: Almost-periodic divisor, almost-periodic function, current of integration, Chern class

Zero distribution for various classes of holomorphic almost-periodic functions in a strip was studied by many authors (cf. [1], [4], [7], [8], [9], [10], [17]). The notion of almost-periodic discrete set appeared in [9] and [17] in connection with these investigations. Its generalization to several complex variables was the notion of almost-periodic divisor, introduced by L. I. Ronkin (cf. [14]) and studied in his works and works of his disciples (cf. [5], [6], [15]). But these notions are not sufficient for a complete description of zero sets of holomorphic almost-periodic functions (cf. [18]): in addition, one needs some topological characteristic, namely, Chern class of the special (generated by an almost-periodic set or a divisor) line bundle over Bohr's compact set (cf. [2], [3]). On the other hand, the class of zero sets of holomorphic functions with almost-periodic modulus in a strip is just the class of almost-periodic discrete sets (cf. [4]). That's why it is natural to obtain a description of zeroes of holomorphic functions with the almost-periodic modulus for several complex variables without using topological terms. This problem is just solved in our paper.

By T_S denote a tube set $\{z = x + iy : x \in \mathbb{R}^m, y \in S\}$, where the base S is a subset of $\mathbb{R}^m$.

Definition 1 . *A continuous function f on T_S is called almost-periodic, if for each sequence $\{f(z + h_n)\}_{h_n \in \mathbb{R}^m}$ of shifts there exists a uniformly convergent on T_S subsequence.*

In particular, for $S = \{0\}$ we obtain the definition of an almost-periodic function on $\mathbb{R}^m$.¹

It follows easily that any almost-periodic function on a tube set with a compact base is bounded.

¹This definition is equivalent to another one that makes use of the notion of an ε -almost period; for $m = 1$ see, for example, [12], the extension to $m > 1$ is trivial.

Definition 2 . Let Ω be a domain in $\mathbb{R}^m$. A continuous function f on T_Ω is called almost-periodic, if its restriction to every tube set T_K with compact base $K \subset \Omega$ is an almost-periodic function on T_K .

Definition 3 (cf. [14], for distributions from $\mathcal{D}'(\mathbb{R})$ cf. also [16]). A distribution $F(z) \in \mathcal{D}'(T_\Omega)$ is called almost-periodic, if for any test-function $\varphi(z) \in \mathcal{D}(T_\Omega)$ the function $\langle F(z), \varphi(z-t) \rangle$ is an almost-periodic function in $t \in \mathbb{R}^m$.

The next assertion is valid.

Theorem 1 (cf. [14]). A distribution $F \in \mathcal{D}'(T_\Omega)$ is almost periodic if and only if for each sequence $\{h^n\} \in \mathbb{R}^m$ there exists a subsequence $\{\tilde{h}^n\}$ such that the sequence of the distributions $F(z + \tilde{h}^n)$ converges uniformly on the sets $\{\kappa(z-t) : t \in \mathbb{R}^m, \kappa \in \mathcal{K}\}$, where $\mathcal{K}$ is any compact family in $\mathcal{D}(T_\Omega)$.

Definition 4 (cf. [14]) The mean value (in the variable $x \in \mathbb{R}^m$) of an almost-periodic distribution F is the distribution $c_F(y) \otimes dx$ with $c_F(y) \in \mathcal{D}'(\Omega)$ and the Lebesgue measure dx on $\mathbb{R}^m$, defined for a test-functions $\varphi \in \mathcal{D}(T_\Omega)$ by the equality

$$\langle c_F(y) \otimes dx, \varphi(z) \rangle = \lim_{N \rightarrow \infty} (2N)^{-m} \int_{\max_j |t_j| < N} \langle F(z), \varphi(z-t) \rangle dt,$$

where $t = (t_1, \dots, t_m) \in \mathbb{R}^m$.

Note that if $F(z)$ is an almost-periodic function on T_Ω , then $c_F(y)$ is a continuous function on Ω . Further, if $F(z)$ is an almost-periodic complex measure on T_Ω , then $c_F(y)$ is a complex measure on Ω as well, and $c_F(y) \otimes dx$ is the weak limit of the measures $F(tx + iy)dx dy$ as $|t| \rightarrow \infty$ (cf. [14]).

By $\mathcal{H}(G)$ denote the space of holomorphic functions on the domain $G \subset \mathbb{C}^m$ with respect to the topology of the uniform convergence on compact subsets of G .

The following assertion is true.

Theorem 2 (cf. [14]). If a function $f \in \mathcal{H}(T_\Omega)$ is almost-periodic, then $\log |f|$ is an almost-periodic distribution on T_Ω .

The main part of the proof of this theorem is the following lemma.

Lemma 1 (cf. [14]). If $f_n \in \mathcal{H}(G)$, $n = 1, 2, \dots$, and $f_n \rightarrow f \not\equiv 0$ in the space $\mathcal{H}(G)$, then $\log |f_n| \rightarrow \log |f|$ in the space $\mathcal{D}'(G)$.

Now let

$$(i/\pi) \partial \bar{\partial} \log |f| = (2/\pi) \sum_{j,k=1}^m \frac{\partial^2 \log |f|}{\partial z_j \partial \bar{z}_k} (i/2) dz_j \wedge d\bar{z}_k. \quad (1)$$

be the current of integration over the divisor d_f of the function $f(z) \in \mathcal{H}(G)$, $z = (z_1, \dots, z_m)$. In the case $m = 1$ this current corresponds to the discrete measure with integer masses equal to the multiplicities of the zeroes of the function f .

Note that all the coefficients of the current (1) are complex measures on G , and the "diagonal" coefficients $\frac{\partial^2 \log |f|}{\partial z_j \partial \bar{z}_j}$ are positive measures (cf. [11]).

Clearly, the differentiation keeps the almost periodicity of distributions. Therefore, it follows from Theorem 2 that all the coefficients of the current (1) are almost-periodic distributions for any holomorphic almost-periodic function on T_Ω . If we replace f by another holomorphic function on T_Ω with the same divisor, then the coefficients of the current (1) do not change. Hence an almost-periodicity of all the coefficients does not imply almost periodicity of the function f itself.

Definition 5 (cf. [5], [6]). *The divisor d_f of a function $f \in \mathcal{H}(T_\Omega)$ is called almost-periodic, if all the coefficients of the current (1) are almost-periodic distributions.*

Note that in [14] a divisor d_f was called almost-periodic, if the measure $\sum_{j=1}^m \frac{\partial^2 \log |f|}{\partial z_j \partial \bar{z}_j}$, was almost-periodic on T_Ω . But that definition is equivalent to the given above (cf. [6]).

There exist almost periodic divisors which cannot be generated by holomorphic almost periodic functions. For example, let $g(w)$ be an entire function on $\mathbb{C}$ with simple zeroes at the points of the standard integer-valued lattice, and let $d[\lambda, \mu]$, $\lambda, \mu \in \mathbb{R}^m$ be the divisor of the function $g(\langle z, \lambda \rangle + i \langle z, \mu \rangle)$. This divisor is periodic for vectors λ, μ that are linearly dependent over $\mathbb{Q}$ or linearly independent over $\mathbb{R}$ (with the periods $\frac{|\mu|^2 \lambda - \langle \lambda, \mu \rangle \mu}{|\lambda|^2 |\mu|^2 - \langle \lambda, \mu \rangle^2}$ and $\frac{|\lambda|^2 \mu - \langle \lambda, \mu \rangle \lambda}{|\lambda|^2 |\mu|^2 - \langle \lambda, \mu \rangle^2}$). Then $d[\lambda, \mu]$ is almost periodic for λ, μ linearly independent over $\mathbb{Q}$ and linearly dependent over $\mathbb{R}$ (for $m = 1$ cf. [18]; since a real linearly transform in $\mathbb{C}^m$ keeps almost-periodicity, the case $m > 1$ follows as well). Besides, the divisor $d[\lambda, \mu]$ for any linearly independent over $\mathbb{Q}$ vectors λ, μ is the divisor of no holomorphic almost periodic function (in the case $m = 1$, i.e., irrational λ/μ cf. [18], for $m > 1$ cf. [15]). A complete description of the divisors of holomorphic almost-periodic functions is contained in the following theorem.

Theorem 3 (for $m = 1$ cf. [2], for $m > 1$ cf. [3]). *A holomorphic bundle over Bohr's compactification K_B of the space $\mathbb{R}^m$ is assigned to each almost-periodic divisor d on a tube domain T_Ω with convex base Ω such that:*

the map $d \mapsto c(d)$, $c(d)$ being the first Chern class of this bundle, is a homomorphism of the semigroup of positive almost-periodic divisors on T_Ω to the cohomology group $H^2(K_B, \mathbb{Z})$, the kernel of this homomorphism is just the set of all divisors of holomorphic almost-periodic functions on T_Ω ,

a finite family $\lambda^j, \mu^j \in \mathbb{R}^m$ corresponds to each cohomology class $c(d)$ such that $c(d) = \sum_j c(d[\lambda^j, \mu^j])$,

the mapping $W : (\lambda, \mu) \mapsto c(d[\lambda, \mu])$ is skew-symmetric and additive in variables $\lambda, \mu \in \mathbb{R}^m$.

A description of zeroes for holomorphic functions of one variable with the almost-periodic modulus is given in the following theorem.

Theorem 4 (cf. [4]; for divisors $d[\lambda, \mu]$, $\lambda, \mu \in \mathbb{R}$ cf. [18]). *A divisor d on a strip is the divisor of some holomorphic function on the strip with almost-periodic modulus if and only if d is almost-periodic.*

Now consider the multidimensional case again. Note that for an almost-periodic divisor d on T_Ω all the coefficients of the current (1) have mean values in x . The imaginary parts of these mean values, i.e., the mean values of the real measures $(2/\pi)\Im \frac{\partial^2 \log |f|}{\partial z_j \partial \bar{z}_k}$ have the form $a_{j,k} dy \otimes dx$, $a_{j,k} \in \mathbb{R}$ (cf. [6]). By $A(d)$ denote the matrix with the entries $a_{j,k}$. In the case $d = d_f$ for an almost-periodic function $f \in \mathcal{H}(T_\Omega)$ we have $A(d) = 0$ (cf. [13]).

Theorem 5 . *A divisor d on a tube domain T_Ω with convex base Ω is the divisor of a holomorphic function with almost-periodic modulus if and only if divisor d is almost-periodic, and the skew-symmetric matrix $A(d)$ is zero.*

To prove this theorem we need the following improvement of Theorem 2.

Theorem 6 . *A function $f \in \mathcal{H}(T_\Omega)$, $f \not\equiv 0$, has almost-periodic modulus if and only if the distribution $\log |f| \in \mathcal{D}'(T_\Omega)$ is almost-periodic.*

Proof of Theorem 6. Let $|f(z)|$ be an almost-periodic function on T_Ω , and let $\{h^n\}$ be an arbitrary sequence from $\mathbb{R}^m$. In order to check that $\log |f|$ is an almost-periodic distribution on T_Ω , we will prove that for any continuous function φ with compact support in T_Ω , the sequence of functions

$$\psi_n(t) = \int \log |f(z + h^n)| \varphi(z - t) dx dy \quad (2)$$

contains a convergent, uniformly on $\mathbb{R}^m$, subsequence. We will prove this assertion by contradiction.

First, since the function $|f(z)|$ is uniformly bounded on T_K for every compact set $K \subset \Omega$, we may assume that the sequence of the functions $\{f(z + h^n)\}$ converges to some function $g(z)$ in the space $\mathcal{H}(T_\Omega)$. Further, since the function $|f(z)|$ is almost-periodic on T_Ω , we may assume that the sequence of the functions $\{|f(z + h^n)|\}$ converges to some function $\Phi(z) \not\equiv 0$ uniformly on each T_K . If the sequence (2) does not converge uniformly on $\mathbb{R}^m$, then for some $\delta > 0$ and some subsequence of n there exist $t^n \in \mathbb{R}^m$ with the property

$$|\psi_n(t^n) - \int \log |g(z)| \varphi(z - t^n) dx dy| \geq \delta. \quad (3)$$

The function $|g(z)| \equiv \Phi(z)$ is almost-periodic on T_Ω , hence we may assume that the same subsequence of the functions $\{|g(z + t^n)|\}$ converges uniformly on each T_K to some function $\Psi(z) \not\equiv 0$. Since the sequence of the functions $\{|f(z + h^n + t)|\}$ converges uniformly in $t \in \mathbb{R}^m$ and $z \in T_K$ to the function $|g(z + t)|$, we see that the subsequence $\{|f(z + h^n + t^n)|\}$ converges to $\Psi(z)$ uniformly on T_K . Also, the subsequences of the functions $\{f(z + h^n + t^n)\}$ and $\{g(z - t^n)\}$ are bounded uniformly on compact subsets of T_Ω , therefore passing to a subsequence again, we get that $f(z + h^n + t^n) \rightarrow H_1(z)$ and $g(z + t^n) \rightarrow H_2(z)$ in the space $\mathcal{H}(T_\Omega)$, and $|H_1(z)| = \Psi(z) = |H_2(z)|$. Using Lemma 1, we obtain that the corresponding subsequences of the functions $\{\log |f(z + h^n + t^n)|\}$ and

$\{\log |g(z + t^n)|\}$ converge, in the sense of distributions, to the same function $\log \Psi(z)$. The last assertion contradicts (3).

On the other hand, let $\log |f(z)|$ be an almost-periodic distribution on T_Ω , and let $\varphi_\varepsilon(z)$ be a nonnegative, depending on $|z|$ smooth function such that $\varphi_\varepsilon(z) = 0$ for $|z| > \varepsilon$ and $\int_{\mathbb{C}^m} \varphi_\varepsilon(z) dx dy = 1$. Evidently, the family of functions $\{\varphi_\varepsilon(z + iy)\}_{|y| \leq C}$ is a compact set in the space $\mathcal{D}(\mathbb{C}^m)$ for every $C < \infty$. Let K be a compact set in Ω and $\varepsilon < \text{dist}\{K, \partial\Omega\}$. Now Theorem 1 implies that the convolution $(\log |f| * \varphi_\varepsilon)(z)$ is an almost-periodic function on T_K . Hence this convolution is bounded on T_K , and the inequality $\log |f(z)| \leq (\log |f| * \varphi_\varepsilon)(z)$ shows that $|f(z)|$ is bounded on T_K as well.

Suppose that $|f|$ is not an almost-periodic function on T_Ω . Then there exists a sequence of functions $\{|f(z + h^n)|\}$, $h^n \in \mathbb{R}^m$, such that every its subsequence does not converge uniformly on $T_{K'}$ for some compact set $K' \subset \Omega$. Without loss of generality it can be assumed that the sequence of functions $\{f(z + h^n)\}$ converges in the space $\mathcal{H}(T_\Omega)$ to some function $g(z)$. It is clear that $g(z)$ is bounded on T_K for every compact set $K \subset \Omega$. Further, by Lemma 1 we get $\log |f(z + h^n)| \rightarrow \log |g(z)|$ in the sense of distributions. Using Theorem 1 and passing to a subsequence, we obtain

$$\int (\log |f(z + h^n)| - \log |g(z)|) \varphi_\varepsilon(z - t - is) dx dy \rightarrow 0. \quad (4)$$

uniformly in $t \in \mathbb{R}^m$ and $s \in K'$. On the other hand, for some $\delta > 0$ and some subsequence of n there exist points $z^n = x^n + iy^n \in T'_K$ such that

$$||f(h^n + x^n + iy^n)| - |g(x^n + iy^n)|| \geq \delta. \quad (5)$$

Passing to a subsequence if necessary, we may assume that $y^n \rightarrow y^0 \in K'$, and the sequences of the functions $\{f(z + h^n + z^n - iy^0)\}$ and $\{g(z + z^n - iy^0)\}$ converge in the space $\mathcal{H}(T_\Omega)$ to functions $H_1(z)$ and $H_2(z)$, respectively. Then Lemma 1 implies that $\log |f(z + h^n + z^n - iy^0)| \rightarrow \log |H_1(z)|$ and $\log |g(z + z^n - iy^0)| \rightarrow \log |H_2(z)|$ in the space $\mathcal{D}'(T_\Omega)$. Taking into account (4), we obtain

$$\int \log |H_1(z)| \varphi_\varepsilon(z - iy^0) dx dy = \int \log |H_2(z)| \varphi_\varepsilon(z - iy^0) dx dy.$$

Since ε is arbitrary small, we get $|H_1(iy^0)| = |H_2(iy^0)|$. At the same time, by (5) we have $|H_1(iy^0)| \neq |H_2(iy^0)|$. This contradiction proves Theorem 6.

Proof of the necessity in Theorem 5. It follows from Theorem 6 that every function $f \in \mathcal{H}(T_\Omega)$ with almost-periodic modulus has an almost-periodic divisor. Further, the mean value $c_{\log |f|}(y) \otimes dx$ of the function $\log |f|$ is the weak limit of the measures $\log |f(tx + iy)| dx \otimes dy$ as $|t| \rightarrow \infty$ in the space $\mathcal{D}'(T_\Omega)$, therefore for all $j \neq k$ the mean values of the distributions

$$\Im \frac{\partial^2 \log |f|}{\partial z_j \partial \bar{z}_k} = \frac{1}{4} \left(\frac{\partial^2}{\partial x_j \partial y_k} - \frac{\partial^2}{\partial x_k \partial y_j} \right) \log |f|$$

equal

$$\begin{aligned} \lim_{|t| \rightarrow \infty} \frac{1}{4} \left(\frac{\partial^2}{\partial x_j \partial y_k} - \frac{\partial^2}{\partial x_k \partial y_j} \right) \log |f(tx + iy)| dx \otimes dy = \\ \frac{1}{4} \left(\frac{\partial^2}{\partial x_j \partial y_k} - \frac{\partial^2}{\partial x_k \partial y_j} \right) c_{\log |f|}(y) \otimes dx = 0. \end{aligned}$$

The necessity of the conditions in Theorem 5 is proved.

The proof of the sufficiency makes use of the following lemmas. As above, $d[\lambda, \mu]$, $\lambda, \mu \in \mathbb{R}^m$ is the divisor of the function $g(\langle z, \lambda \rangle + i\langle z, \mu \rangle)$, where $g(w)$ is an entire function on $\mathbb{C}$ with simple zeroes at the points of the standard integer-valued lattice.

Lemma 2 . *The divisor $d[\lambda, \mu]$ with $\lambda = t\mu$, $\lambda \in \mathbb{R}^m$, $t \in \mathbb{R}$, is the divisor of an entire function on $\mathbb{C}^m$ with almost-periodic modulus.*

Proof of Lemma 2. After a suitable regular real linear transform we obtain the case $\mu = (1, 0, \dots, 0)$, i.e., the case of a divisor depending only on one coordinate, therefore the assertion of our lemma is a consequence of Theorem 4.

Further, let $e^1, \dots, e^m$ be the coordinate vectors in $\mathbb{C}^m$.

Lemma 3 . *The entries $a_{j,k}$ of the matrix $A_0 = A(d[e^1, e^2])$ vanish for $(j, k) \neq (1, 2)$ or $(2, 1)$, and $a_{1,2} = -1$, $a_{2,1} = 1$.*

Proof of Lemma 3. The divisor of the function $g(z_1 + iz_2)$ does not depend on variables z_j with $j > 2$, hence the distributions $\Im \frac{\partial^2 \log |g(z_1 + iz_2)|}{\partial z_j \partial \bar{z}_k}$ vanish for $(j, k) \neq (1, 2)$ or $(2, 1)$.

Consider the expression

$$(L_z \log |g(z_1 + iz_2)|, \varphi(z_1 + t_1, z_2 + t_2)), \quad (t_1, t_2) \in \mathbb{R}^2, \quad (6)$$

for $L_z = \frac{2}{\pi} \Im \frac{\partial^2}{\partial \bar{z}_1 \partial z_2}$ and a function $\varphi(z) \geq 0$ from the space $\mathcal{D}(\mathbb{C}^2)$. In the coordinates $\zeta_1 = z_1 + iz_2$, $\zeta_2 = z_1 - iz_2$, it has a form

$$\frac{1}{4} (\tilde{L}_\zeta \log |g(\zeta_1)|, \varphi((\zeta_1 + \zeta_2)/2 + t_1, (\zeta_1 - \zeta_2)/2i + t_2))$$

with

$$\tilde{L}_\zeta = \frac{2}{\pi} \Re \left(\frac{\partial^2}{\partial \zeta_1 \partial \bar{\zeta}_1} - \frac{\partial^2}{\partial \zeta_2 \partial \bar{\zeta}_1} + \frac{\partial^2}{\partial \zeta_1 \partial \bar{\zeta}_2} - \frac{\partial^2}{\partial \zeta_2 \partial \bar{\zeta}_2} \right).$$

Using the definition of g and properties of the Laplace operator, we get

$$\tilde{L}_\zeta \log |g(\zeta_1)| = \frac{2}{\pi} \frac{\partial^2}{\partial \zeta_1 \partial \bar{\zeta}_1} \log |g(\zeta_1)| = \sum_{q_1, q_2 \in \mathbb{Z}} \delta(\zeta_1 - q_1 - iq_2) \otimes d\xi d\eta,$$

where δ is the Dirac function on the plane, $\xi = \Re \zeta_2$, $\eta = \Im \zeta_2$. Therefore, (6) is equal to

$$\frac{1}{4} \sum_{q_1, q_2 \in \mathbb{Z}} \int_{\mathbb{C}} \varphi(t_1 + (q_1 + iq_2 + \xi + i\eta)/2, t_2 + (q_1 + iq_2 - \xi - i\eta)/2i) d\xi d\eta.$$

Substituting $\xi - q_1$, $\eta + q_2$ for u , v , respectively, we get

$$\frac{1}{4} \sum_{q_1, q_2 \in \mathbb{Z}} \int_{\mathbb{C}} \varphi(u/2 + iv/2 + t_1 + q_1, -v/2 + iu/2 + t_2 + q_2) du dv. \quad (7)$$

Since the divisor d_{e^1, e^2} has period 1 in each variable, we see that the mean value of (6) is the integral of (7) over the square $0 \leq t_1 \leq 1, 0 \leq t_2 \leq 1$. Then it is equal to the integral

$$\frac{1}{4} \int_{\mathbb{R}^4} \varphi(u/2 + iv/2 + t_1, -v/2 + iu/2 + t_2) du dv dt_1 dt_2.$$

Finally, substituting $u/2 + t_1 = x_1, v/2 = y_1, t_2 - v/2 = x_2, u/2 = y_2$, we obtain the equality

$$\begin{aligned} \int_0^1 \int_0^1 (L_z \log |g(z_1 + iz_2)|, \varphi(z_1 + t_1, z_2 + t_2)) dt_1 dt_2 = \\ \int_{\mathbb{R}^4} \varphi(x_1 + iy_1, x_2 + iy_2) dx_1 dy_1 dx_2 dy_2, \end{aligned}$$

hence the mean value of the distribution $L_z \log |g(z_1 + iz_2)|$ is the Lebesgue measure in $\mathbb{C}^2$. The lemma is proved.

By (λ, μ) denote the matrix product $(\lambda_j \mu_k)_{j,k=1}^m$ of the vectors $\lambda = (\lambda_1, \dots, \lambda_m), \mu = (\mu_1, \dots, \mu_m) \in \mathbb{R}^m$.

Lemma 4 . *For any $\lambda, \mu \in \mathbb{R}^m$, the matrix $A(d[\lambda, \mu])$ equals the difference $(\mu, \lambda) - (\lambda, \mu)$.*

Proof of Lemma 4. If λ, μ are linearly dependent over $\mathbb{R}$, then $(\mu, \lambda) - (\lambda, \mu) = 0$. On the other hand, it follows from Lemma 2 that in this case the divisor $d[\lambda, \mu]$ is the divisor of some holomorphic in $\mathbb{C}^m$ function with almost-periodic modulus. Using the proved part of Theorem 5, we have $A(d[\lambda, \mu]) = 0$.

Let λ, μ be linearly independent over $\mathbb{R}$. The divisor $d[\lambda, \mu]$ is the divisor $d[e^1, e^2]$ in the coordinates $\zeta = Bz$ for some real nondegenerate matrix B with the first and second rows λ and μ , respectively. Note that the matrix $A(d)$ is the matrix of the mean values for the matrix $\frac{1}{2i}(D(z) - \bar{D}(z))$, where

$$D(z) = \left(\frac{\partial^2 \log |g(\langle z, \lambda \rangle + i \langle z, \mu \rangle)|}{\partial z_j \partial \bar{z}_k} \right),$$

$\bar{D}$ being the matrix with all the entries complex conjugated to the corresponding entries of the matrix D . Therefore $D(z) = B' D(\zeta) B$, B' being the transpose matrix to B , and $A(d[\lambda, \mu]) = B' A_0 B$ for the matrix A_0 from the previous lemma. This completes the proof of Lemma 4.

Lemma 5 . *If numbers $\alpha_j, \beta_j \in \mathbb{R}, j = 1, \dots, n$, satisfy the condition $\sum_1^n \alpha_j \beta_j = 0$, then for some $\gamma_k \in \mathbb{R}, \nu^k \in \mathbb{R}^m, k = 1, \dots, N$, we get*

$$\sum_1^n W(\alpha_j e^1, \beta_j e^2) = \sum_1^N W(\gamma_k \nu^k, \nu^k), \quad (8)$$

W being the mapping from Theorem 3.

Proof of Lemma 5. The case $n = 1$ means that the left-hand side of (8) vanishes. For $n > 1$ we have

$$\begin{aligned} W(\alpha_{n-1}e^1, \beta_{n-1}e^2) + W(\alpha_n e^1, \beta_n e^2) &= W(\alpha_{n-1}e^1, \alpha_n e^1) + W(\beta_n e^2, \beta_n \alpha_n / \alpha_{n-1} e^2) \\ &+ W(\alpha_n e^1 + \alpha_n \beta_n / \alpha_{n-1} e^2, \alpha_{n-1} e^1) + \beta_n e^2 + W(\alpha_{n-1} e^1, (\beta_{n-1} + \beta_n \alpha_n / \alpha_{n-1}) e^2). \end{aligned}$$

The first three terms of the right-hand side have the form $W(\gamma\nu, \nu)$, $\gamma \in \mathbb{R}, \nu \in \mathbb{R}^m$. Subtracting these terms from the left-hand side of (8), we get

$$\sum_1^{n-2} W(\alpha_j e^1, \beta_j e^2) + W(\alpha_{n-1} e^1, (\beta_{n-1} + \beta_n \alpha_n / \alpha_{n-1}) e^2).$$

Hence the lemma can be proved by induction over n .

Lemma 6 . Let vectors $\lambda^j, \mu^j \in \mathbb{R}^m$, $j = 1, \dots, n$ be such that the matrix $\sum_1^n (\lambda^j, \mu^j)$ is symmetric. Then

$$\sum_1^n W(\lambda^j, \mu^j) = \sum_1^N W(\gamma_k \nu^k, \nu^k) \quad (9)$$

for some $\gamma_k \in \mathbb{R}$, $\nu^k \in \mathbb{R}^m$, $k = 1, \dots, N$.

Proof of Lemma 6. The vectors λ^j, μ^j are linear combinations of the vectors $e^1, \dots, e^m$, therefore the left-hand side of (9) has the form

$$\sum_{1 \leq p, q \leq m} \left(\sum_{j=1}^{M(p,q)} W(\alpha_{j,p} e^p, \beta_{j,q} e^q) \right) \quad (10)$$

with $\alpha_{j,p}, \beta_{j,q} \in \mathbb{R}$. The mapping W is skew-symmetric, hence we may assume that all the terms in (10) vanish for $p > q$, and the entries of the corresponding matrix $\left(\sum_{j=1}^{M(p,q)} \alpha_{j,p} \beta_{j,q} \right)_{p,q=1}^m$ vanish for all $p > q$. Since this matrix coincides with the symmetric matrix $\sum_1^n (\lambda^j, \mu^j)$, we see that $\sum_{j=1}^{M(p,q)} \alpha_{j,p} \beta_{j,q} = 0$ for $p < q$ as well. Now it follows from Lemma 5 that for $p < q$ the sum

$$\sum_{j=1}^{M(p,q)} W(\alpha_{j,p} e^p, \beta_{j,q} e^q)$$

has the form of the right-hand side of (9). The terms of (10) with $p = q$ have already the form $W(\gamma\nu, \nu)$. The lemma is proved.

Proof of the sufficiency in Theorem 5. Let d be a divisor in T_Ω such that $A(d) = 0$. It follows from Theorem 3 that there exist $\lambda^j, \mu^j \in \mathbb{R}^m$, $j = 1, \dots, n$, such that the sum $d + \sum_j d[\lambda^j, \mu^j]$ is the divisor of a holomorphic almost-periodic function. Now, by [13], $A(d + \sum_1^n d[\lambda^j, \mu^j]) = 0$. Since the mapping $d \mapsto A(d)$ is a homomorphism,

we get $\sum_1^n (\lambda^j, \mu^j) - (\mu^j, \lambda^j) = \sum_1^n A(d[\lambda^j, \mu^j]) = 0$, i.e., the matrix $\sum_1^n (\lambda^j, \mu^j)$ is symmetric. Using Lemma 6, we get (9) for some $\gamma_k \in \mathbb{R}$, $\nu^k \in \mathbb{R}^m$, $k = 1, \dots, N$. Therefore,

$$\begin{aligned} c(d + \sum_1^N d[\gamma_k \nu^k, \nu^k]) &= c(d) + \sum_1^N W(\gamma_k \nu^k, \nu^k) \\ &= c(d) + \sum_1^n W(\lambda^j, \mu^j) = c(d + \sum_1^n d[\lambda^j, \mu^j]) = 0. \end{aligned}$$

An application of Theorem 3 yields that there exists an almost-periodic function $F \in \mathcal{H}(T_\Omega)$ with the divisor $d + \sum_1^N d[\gamma_k \nu^k, \nu^k]$. Using Lemma 2, we can take functions $f_k \in \mathcal{H}(T_\Omega)$ with the divisors $d[\gamma_k \nu^k, \nu^k]$ and almost-periodic modula. The function $f(z) = F(z)(\prod_1^N f_k(z))^{-1}$ is holomorphic on T_Ω and has the divisor d . Then Theorem 6 implies that the distributions $\log |F|$ and $\log |f_k|$, $k = 1, \dots, N$, are almost-periodic. Hence the distribution $\log |f| = \log |F| - \sum_1^N \log |f_k|$ is almost-periodic as well. Using Theorem 6 again, we see that the function $|f|$ is almost-periodic. This completes the proof of Theorem 5.

References

- [1] H. Bohr, *Zur Theorie der Fastperiodischen Funktionen, III Teil; Dirichletentwicklung Analytischer Funktionen*, Acta math. **47** (1926), 237-281.
- [2] Favorov, S.Yu. *Zeros of holomorphic almost periodic functions*, Journal d'Analyse Mathematique 84, (2001), 51-66.
- [3] Favorov, S.Yu., *Almost periodic divisors, holomorphic functions, and holomorphic mappings*, Bull.Sci.Math. 127, (2003), 859-883.
- [4] Favorov, S.Yu., Rashkovskii, A.Ju., and Ronkin, L.I., *Almost periodic divisors in a strip*, Journal d'Analyse Math. 74, (1998), 325-345.
- [5] Favorov, S.Yu., Rashkovskii, A.Ju., and Ronkin, L.I., *Almost periodic currents and holomorphic chains*, C. R. Acad. Sci. Paris 327, Serie I (1998), 302-307.
- [6] Favorov, S.Yu., Rashkovskii, A.Ju., and Ronkin, L.I., *Almost periodic currents, divisors and holomorphic chains in tube domains*, Israel. Math. Conf. Proc. 15, (2001), 67-88.
- [7] B. Jessen, *Über die Nullstellen einer analitischen fastperiodischen Funktions, Eine Verallgemeinerung der Jensenschen Formel*, Math. Ann. **108** (1933), 485-516.
- [8] B. Jessen, H. Tornehave, *Mean motions and zeros of almost periodic functions*, Acta Math. **77** (1945), 137-279.
- [9] Krejn, M.G., and Levin, B.Ja., *On almost periodic functions of exponential type*, Dokl. AN SSSR **64** (1949), no. 3, 285-287 (Russian).

- [10] Levin, B. Ja., *Distribution of Zeros of Entire Functions*. Transl. of Math. Monograph, Vol. 5, AMS Providence, no. 1, MR 19, 1964, 403p.
- [11] Lelong, P., and Gruman, L., *Functions of Several Complex Variables*, Springer-Verlag. Berlin-Heidelberg, 1986.
- [12] Levitan, B. M., *Almost periodic functions*. - M.: Gostehizdat, 1953, 396 p, (Russian).
- [13] Ronkin, L. I., *Jessen's theorem for holomorphic almost periodic mappings*, Ukrainsk. Mat. Zh. **42** (1990), 1094-1107 (Russian).
- [14] Ronkin, L. I., *Almost periodic distributions and divisors in tube domains*, Zap. Nauchn. Sem. POMI **247** (1997), 210-236 (Russian).
- [15] Ronkin, L. I., *Almost periodic divisors and the spatial extension of almost periodic measures*, Linear Topological Spaces and Complex Analysis 3, (1997), 152-156.
- [16] Schwartz, L., *Théorie des distributions 1*. Hermann, Paris, 1950. – 350p.
- [17] Tornehave, H., *Systems of zeros of holomorphic almost periodic functions*, Kobenhavns Universitet Matematisk Institut, Preprint no. 30, 1988.
- [18] Tornehave, H., *On the zeros of entire almost periodic functions*, The Harald Bohr Centenary (Copenhagen 1987). Math. Fys. Medd. Danske 42 (3), (1989), 125-142.

Department of Mathematics
 Kharkov National University
 Svobody sq., 4, Kharkov 61077
 Ukraine
 e-mail: favorov@assa.kharkov.ua